

# Nontrivial Fermion States in Supersymmetric Minisuperspace\*

András Csordás<sup>†</sup> and Robert Graham

Fachbereich Physik, Universität-Gesamthochschule Essen  
45117 Essen  
Germany

## Abstract

In this lecture supersymmetric minisuperspace models of any Bianchi type within class A of the classification of Ellis and McCallum are considered. The algebra of the supersymmetry generators, the Lorentz generators, the diffeomorphism generators and the Hamiltonian generator is determined explicitly and found to close. Different from earlier work it is established that physical states, which are annihilated by all these generators, exist in *all* sectors of these models with fixed even fermion number. A state in the 4-fermion sector of the Bianchi type IX model is considered as a specific example, which satisfies the ‘no-boundary’ condition of Hartle and Hawking. The conclusion is that supersymmetric minisuperspace models have a much richer manifold of physical states than had been recognized before.

While minisuperspace models have long served as a useful laboratory for new ideas in quantum gravity, it was only rather recently that also the quantum theory of *supersymmetric* minisuperspace models has attracted a broad interest (see e.g. [1]-[13]). It was found [4] that supersymmetric versions have special explicit analytical solutions in the *empty* and *filled* fermion sectors which can be interpreted as wormhole states [6] (and with a different homogeneity condition for the gravitino, as Hartle-Hawking no-boundary states [13]). However, even more recently it has been shown by a simple scaling argument, that the special solutions found in the minisuperspace models have no direct counterpart in

---

\*Invited lecture presented at the ‘First Mexican School on Gravitation and Mathematical Physics’,

Guanajuato, GTO., Mexico, Dec. 12–16, 1994

<sup>†</sup>Permanent address: Research Institute for Solid State Physics, P.O. Box 49, H1525 Budapest, Hungary

4-dimensional supergravity, because there states in the empty (and also in the filled) sector cannot exist [14]. In this light another result found some time ago [5] and independently confirmed by several authors [6, 7, 8, 9] appears very puzzling: namely that the solutions in the empty and filled fermion sectors are the *only* physical states satisfying all constraints for all supersymmetric minisuperspace models of Bianchi type in class A [15] (without matter-coupling and with the exception, with a certain operator ordering [8, 9], of Bianchi type I). For anisotropic supersymmetric minisuperspace models with a non-vanishing cosmological constant [10, 11, 12] it even is reported in the literature that *no* physical states exist, at all. Taken together these results imply that the physical states found in the minisuperspace models have no counterpart in the full theory, and vice-versa, which would render supersymmetric minisuperspace models useless as models of full supergravity, contrary to the situation in pure gravity. So, why this lecture?

Because there is a paradox: How can the supersymmetric models be more constrained than the corresponding models in pure gravity, even though supergravity certainly has *more* physical degrees of freedom than gravity, not less (namely, in full supergravity, per space-point there are two additional physical degrees of freedom of the gravitino, which corresponds to two additional degrees of freedom in the minisuperspace models).

The work [19] discussed in this lecture has the purpose to clarify these issues for the supersymmetric models of Bianchi type within class A of the classification of Ellis and McCallum [15], restricting ourselves to the case of a vanishing cosmological constant.

The starting point is the Lagrangean of  $N = 1$  supergravity given in [16]. We adopt all spinor conventions given there and shall also make free use of the excellent account of the Hamiltonian form of  $N = 1$  supergravity in the metric representation given in [17]. In the metric representation, which we shall use, the independent variables are taken to be the tetrad components  $e_p^a$  (with Einstein indices  $p = 1, 2, 3$  from the middle of the alphabet and Lorentz indices  $a = 0, 1, 2, 3$  from the beginning of the alphabet) and the Grassmannian components  $\psi_p^\alpha$  and their adjoints  $\bar{\psi}_p^{\dot{\alpha}}$  (with spinor index  $\alpha = 1, 2$ ;  $\dot{\alpha} = 1, 2$ ) of the Rarita-Schwinger field. The  $e_p^a$  form the metric tensor  $h_{pq} = e_p^a e_{qa}$  on the space-like homogeneity 3-surfaces in the symmetric basis of 1-forms  $\omega^p$ , satisfying in Bianchi type models of class A

$$d\omega^p = \frac{1}{2} \frac{m^{pq}}{h^{1/2}} \epsilon_{qrs} \omega^r \wedge \omega^s \quad (1)$$

Here  $h = \det h_{pq}$  and  $\epsilon_{qrs}$  denotes the components of the Levi-Civita tensor and we shall use the notation  $V = \int \omega^1 \wedge \omega^2 \wedge \omega^3$ . The constant symmetric matrix  $m^{pq}$  is fixed by the chosen Bianchi type [18]. It is invariant and transforms as a tensor under all coordinate changes from one *symmetric* basis to another one. Due to the choice of a symmetric basis the  $e_p^a$  and  $\psi_p^\alpha$ ,  $\bar{\psi}_p^{\dot{\alpha}}$  are functions of time only.

From the Lagrangean one defines, as usual, the generalized momenta  $\hat{p}_a^p$  and  $\hat{\pi}_\alpha^p$ ,  $\hat{\pi}_{\dot{\alpha}}^p$  of  $e_p^a$  and  $\psi_p^\alpha$ ,  $\bar{\psi}_p^{\dot{\alpha}}$  respectively. The Poisson brackets must be replaced by Dirac brackets due to the appearance and subsequent elimination of second class constraints, by which the adjoint quantities  $\bar{\psi}_p^{\dot{\alpha}}$ ,  $\hat{\pi}_{\dot{\alpha}}^p$  are eliminated. The Dirac brackets are decoupled by the introduction of new  $p_+^p{}_a$  and  $\pi^p{}_\alpha$ .

The only non-vanishing Dirac brackets then are

$$\begin{aligned}\{e_p^a, p_+^q{}_b\}^* &= \delta_p^q \delta_b^a \\ \{\pi_p^\alpha, \psi_p^q\}^* &= -\delta_p^q \delta_\beta^\alpha.\end{aligned}\quad (2)$$

The supersymmetry generators  $S_\alpha$ ,  $\bar{S}_{\dot{\alpha}}$  and Lorentz generators  $J_{\alpha\beta}$ ,  $\bar{J}_{\dot{\alpha}\dot{\beta}}$  in this representation are obtained as

$$\begin{aligned}S_\alpha &= -\mathcal{C}_{pr}^{\dot{\alpha}\beta} \left( \frac{1}{2} V m^{pq} e_q^a + \frac{i}{2} p_+^{pa} \right) \sigma_{a\alpha\dot{\alpha}} \pi^r{}_\beta \\ \bar{S}_{\dot{\alpha}} &= \left( \frac{1}{2} V m^{pq} e_q^a - \frac{i}{2} p_+^{pa} \right) \sigma_{a\alpha\dot{\alpha}} \psi_p^\alpha\end{aligned}\quad (3)$$

and

$$\begin{aligned}J_{\alpha\beta} &= +\frac{1}{2}(\sigma^{ac}\epsilon)_{\alpha\beta} (e_{pa} p_+^p{}_c - e_{pc} p_+^p{}_a) \\ &\quad -\frac{1}{2}(\psi_{p\alpha} \pi^p{}_\beta + \psi_{p\beta} \pi^p{}_\alpha) \\ \bar{J}_{\dot{\alpha}\dot{\beta}} &= -\frac{1}{2}(\bar{\epsilon}^{ac})_{\dot{\alpha}\dot{\beta}} (e_{pa} p_+^p{}_c - e_{pc} p_+^p{}_a).\end{aligned}\quad (4)$$

with

$$\mathcal{C}_{pq}^{\dot{\alpha}\alpha} = -\frac{1}{2Vh^{1/2}} (i\hbar_{pq} n^a - \varepsilon_{pqr} e^{ra}) \bar{\sigma}_a^{\dot{\alpha}\alpha}.\quad (5)$$

Canonical quantization is achieved by putting

$$p_+^p{}_a = -i\hbar(\partial/\partial e_p^a) \quad \pi^p{}_\alpha = -i\hbar(\partial/\partial \psi_p^\alpha).\quad (6)$$

In  $S_\alpha$  there is then an ordering ambiguity between  $p_+^p{}_a$  and  $\mathcal{C}_{pr}^{\dot{\alpha}\beta}$ , which is resolved by adopting the ordering as written in eq. (3) [20]. Now a somewhat lengthy calculation gives the following generator algebra [19, 20]: The commutators of  $J_{\alpha\beta}$ ,  $\bar{J}_{\dot{\alpha}\dot{\beta}}$  with any operator is determined by its changes under infinitesimal Lorentz transformations. The remaining commutators and anti-commutators are

$$[S_\alpha, S_\beta]_+ = 0 = [\bar{S}_{\dot{\alpha}}, \bar{S}_{\dot{\beta}}]_+ \quad (7)$$

$$[S_\alpha, \bar{S}_{\dot{\alpha}}]_+ = -\frac{\hbar}{2} H_{\alpha\dot{\alpha}} \quad (8)$$

$$[H_{\alpha\dot{\alpha}}, S_\beta]_- = -i\hbar \varepsilon_{\alpha\beta} \bar{D}_{\dot{\alpha}}^{\dot{\beta}\dot{\gamma}} \bar{J}_{\dot{\beta}\dot{\gamma}} \quad (9)$$

$$\begin{aligned}[H_{\alpha\dot{\alpha}}, \bar{S}_{\dot{\beta}}]_- &= i\hbar \varepsilon_{\dot{\alpha}\dot{\beta}} J_{\beta\gamma} D_\alpha^{\beta\gamma} \\ &= i\hbar \varepsilon_{\dot{\alpha}\dot{\beta}} \left[ D_\alpha^{\beta\gamma} J_{\beta\gamma} + i\hbar \bar{E}_\alpha^{\dot{\gamma}\dot{\delta}} \bar{J}_{\dot{\gamma}\dot{\delta}} + \frac{i\hbar n^a}{Vh^{1/2}} \sigma_{a\alpha\dot{\gamma}} \bar{S}^{\dot{\gamma}} \right].\end{aligned}\quad (10)$$

The operator  $H_{\alpha\dot{\alpha}}$  is here *defined* by the anticommutator (8). We checked that it differs from the diffeomorphism — and Hamiltonian constraints  $\tilde{H}_{\alpha\dot{\alpha}} = \sigma_{a\alpha\dot{\alpha}}(e_p^a H^p + n^a H)$  only by terms proportional to Lorentz generators. The commutators  $[\bar{S}_{\dot{\beta}}, H_{\alpha\dot{\alpha}}]_-$  and  $[S_{\beta}, H_{\alpha\dot{\alpha}}]_-$  are the essential new results which will turn out to be crucial, in the following.  $D_{\alpha}^{\beta\gamma}$ ,  $\bar{E}_{\dot{\alpha}}^{\dot{\gamma}\dot{\delta}}$  are Grassmann-odd operators whose explicit form we shall not need. Eqs. (7)-(10) demonstrate explicitly that the algebra of the generators closes, and thereby confirm the assumption made in earlier work on the same minisuperspace models.

The physical states of these models are given by all the states which are annihilated by the generators  $S_{\alpha}$ ,  $\bar{S}_{\dot{\alpha}}$ ,  $J_{\alpha\beta}$ ,  $\bar{J}_{\dot{\alpha}\dot{\beta}}$ ,  $H_{\alpha\dot{\alpha}}$ . The Lorentz generators automatically annihilate all states which are Lorentz scalars. Therefore, it is sufficient to demand that physical states are Lorentz scalars and annihilated by  $S_{\alpha}$  and  $\bar{S}_{\dot{\alpha}}$ ; their annihilation by  $H_{\alpha\dot{\alpha}}$  is then automatically guaranteed by the generator algebra. The form of the constraint operators guarantees that physical states have a fixed fermion number  $F = \psi_p^{\alpha} \partial / \partial \psi_p^{\alpha}$ , which must be an even number in Lorentz-invariant states and ranges from 0 to 6 in the present models.

The physical states in the sectors  $F = 0$  and  $F = 6$  are easily obtained, and are, respectively, given by [20]

$$\begin{aligned}\Psi_0 &= \text{const } e^{\frac{V}{2\hbar} m^{pq} h_{pq}} \\ \Psi_6 &= \text{const } e^{-\frac{V}{2\hbar} m^{pq} h_{pq}} \prod_r (\psi_r)^2\end{aligned}\tag{11}$$

reproducing a well known result [4]-[9].

In order to show that there exist physical states in the 2-fermion sector as well we consider the wave-function [19]

$$\Psi_2 = \bar{S}_{\dot{\alpha}} \bar{S}^{\dot{\alpha}} f(h_{pq}),\tag{12}$$

with  $\bar{S}^{\dot{\alpha}} f \neq 0$ , where  $f$  is a function of the  $h_{pq}$  only, and therefore, like  $\bar{S}_{\dot{\alpha}} \bar{S}^{\dot{\alpha}}$ , a Lorentz scalar. Therefore  $\Psi_2$  *automatically* satisfies the Lorentz constraints and the  $\bar{S}$ -constraints because of eq. (7). The only remaining condition is  $S_{\alpha} \Psi_2 = 0$ , which, after the use of eqs. (12), (8), reduces to

$$[H_{\alpha\dot{\alpha}}, \bar{S}^{\dot{\alpha}}]_- f + 2\bar{S}^{\dot{\alpha}} H_{\alpha\dot{\alpha}} f = 0\tag{13}$$

The first term, thanks to eq. (10), reduces to terms proportional to  $J_{\beta\gamma}$ ,  $\bar{J}_{\dot{\gamma}\dot{\delta}}$ ,  $\bar{S}^{\dot{\gamma}}$ . The terms with  $J_{\beta\gamma}$ ,  $\bar{J}_{\dot{\gamma}\dot{\delta}}$  vanish because  $f$  is a Lorentz scalar. In the remainder  $\bar{S}^{\dot{\gamma}}$  can be factored out to the left, using  $[\bar{S}^{\dot{\gamma}}, \sigma_{a\alpha\dot{\gamma}} n^a / h^{1/2}]_- = 0$ . Then it is seen that eq. (13) is solved if  $f$  satisfies the Wheeler-DeWitt equation [20]

$$\left( H_{\alpha\dot{\alpha}}^{(0)} - \frac{\hbar^2}{V h^{1/2}} n^a \sigma_{a\alpha\dot{\alpha}} \right) f(h_{pq}) = 0\tag{14}$$

where  $H_{\alpha\dot{\alpha}}^{(0)}$  consists only of the bosonic terms of  $H_{\alpha\dot{\alpha}}$ , i.e. of the terms which remain if  $\pi^p_\alpha$  is brought to the right and then equated to zero. Any solution of this Wheeler-DeWitt equation, which may be specified further by imposing boundary conditions, determines a solution in the 2-fermion sector via eq. (12), with a definite dependence on the fermionic variables. It should be noted that the norm of  $\Psi_2$  vanishes due to the appearance of  $\bar{S}_{\dot{\alpha}}$  in eq. (12) and the fact that  $S_\alpha$  is the adjoint of  $\bar{S}_{\dot{\alpha}}$ . However, including the necessary gauge-fixing condition in the measure of the scalar product, the norm of  $\Psi_2$  (and of  $\Psi_4$  to be considered below) will not vanish.

We now turn to physical states in the 4-fermion sector. Similarly to eq. (12) the wave-function

$$\Psi_4 = S^\alpha S_\alpha g(h_{pq}) \prod_{r=1}^3 (\psi_r)^2. \quad (15)$$

with  $\Psi_4 \neq 0$  automatically satisfies the Lorentz constraints and the  $S_\alpha$ -constraint. It remains to satisfy the  $\bar{S}_{\dot{\alpha}}$  constraint. A calculation similar to the one leading from eq. (12) to eq. (15) then gives [19, 20] the Wheeler-DeWitt equation.

$$H_{\alpha\dot{\alpha}}^{(1)} g(h_{pq}) = 0 \quad (16)$$

where  $H_{\alpha\dot{\alpha}}^{(1)}$  consists of those terms of  $H_{\alpha\dot{\alpha}}$  which remain if the  $\pi^p_a$  are brought to the left and then equated to zero. Eq. (16) is slightly different from that obeyed by the amplitude in the 2-fermion sector, but the degree of generality of the solution is the same.

Our results differ from earlier work on the supersymmetric Bianchi models in class A [5]-[9] which concluded that physical states in the 2- and 4-fermion sectors do not exist. In fact the ansatz made there was overly restrictive and did not include all the Lorentz invariants contained in the states (12) and (15).

What about the relation to full supergravity? Provided the algebra of the local generators of the constraints still has a form like eqs. (7)-(10) physical states still exist which look somewhat like  $\Psi_2$  and  $\Psi_4$ , but contain formal products of  $(\bar{S})^2$  or  $(S)^2$  over all points of the space-like 3-surface, thus leading to states with infinite fermion number. Such states have recently been discussed in [14]. The new physical states discussed in the present paper are the direct minisuperspace analogues of such states in full supergravity; even though, due to the reduction to minisuperspace the fermion number is, of course, finite. The new states display the same richness of gravitational dynamical behavior as the Bianchi models in pure gravity. In this respect they differ qualitatively from the earlier found states in the empty and the full fermion sectors, which are highly symmetric and do not describe the very rich dynamical behavior of the Bianchi models in the classical limit. The new physical states described here span an infinite-dimensional Hilbert space just as the states of the Bianchi models of pure gravity. Just how this Hilbert space ought to be constructed remains one of the interesting open questions.

Before ending let us consider a specific example, namely the case of Bianchi type IX. There the metric  $h_{pq}$  can be parametrized by the Euler angles  $\varphi, \vartheta, \chi$  contained in a rotation matrix  $\Omega_{pi}$  and three scale parameters

$$(e^{2\beta})_{ij} = e^{2\alpha} \text{diag} (e^{2\beta_+ + 2\sqrt{3}\beta_-}, e^{2\beta_+ - 2\sqrt{3}\beta_-}, e^{-4\beta_+}) \quad (17)$$

via  $h_{pq} = \Omega_{pi}(e^{2\beta})_{ij}\Omega_{qj}$ .

Let us consider, in particular, eq.(16) for the 4-fermion sector. The three equations

$e_p^a \bar{\sigma}_a^{\dot{\alpha}\alpha} H_{\alpha\dot{\alpha}}^{(1)} g = 0$  turn out to be automatically satisfied if  $g$  is *independent* of the Euler angles  $\varphi, \vartheta, \chi$ . The only remaining constraint  $n^a \bar{\sigma}_a^{\dot{\alpha}\alpha} H_{\alpha\dot{\alpha}}^{(1)} g = 0$  then takes the form

$$\left[ -\frac{\hbar^2}{V^2} \left( \frac{\partial}{\partial \alpha} \right)^2 + \frac{\hbar^2}{V^2} \left( \frac{\partial}{\partial \beta_+} \right)^2 + \frac{\hbar^2}{V^2} \left( \frac{\partial}{\partial \beta_-} \right)^2 + \left( \frac{\partial \phi}{\partial \alpha} \right)^2 - \left( \frac{\partial \phi}{\partial \beta_+} \right)^2 - \left( \frac{\partial \phi}{\partial \beta_-} \right)^2 + \frac{\hbar}{V} \left( -\frac{\partial^2 \phi}{\partial \alpha^2} + \frac{\partial^2 \phi}{\partial \beta_+^2} + \frac{\partial^2 \phi}{\partial \beta_-^2} \right) \right] g(\alpha, \beta_+, \beta_-) = 0 \quad (18)$$

with

$$\phi = \frac{1}{2} e^{2\alpha} (2e^{2\beta_+} \cosh 2\sqrt{3}\beta_- + e^{-4\beta_+}) .$$

This equation is of exactly the same form as given in ref. [4]. Even though  $g = \exp(-V\phi/\hbar)$  solves eq. (18) exactly [4], this solution is not permitted here, because it would fail to satisfy the condition  $\Psi_4 \neq 0$ . Remarkably, however, *another* equally simple exact solution of eq. (18) exists without this defect. It is

$$g = \text{const} \exp \left[ -\frac{V}{2\hbar} e^{2\alpha} (2e^{2\beta_+} (\cosh 2\sqrt{3}\beta_- - 1) + e^{-4\beta_+} - 4e^{-\beta_+} \cosh \sqrt{3}\beta_-) \right] . \quad (19)$$

As has been discussed elsewhere (see e.g. the last reference [4]), this state satisfies the requirements of a Hartle-Hawking ‘no-boundary’ state. In Bianchi type IX models with the homogeneity condition  $\psi^\alpha = \psi_p^\alpha(t) \omega^p$  such a state, up til now, was mysteriously absent [6] which led to the proposal to modify the homogeneity condition for the Rarita-Schwinger field [13]. It is therefore very interesting and gratifying to see now such a state appear among the new states (15) without any modification in the homogeneity condition.

In summary this work demonstrates that anisotropic supersymmetric minisuperspace models have a much richer manifold of physical states than had been recognized up to now with clear correspondences with states both in minisuperspace models of pure gravity and in full supergravity.

This work has been supported by the Deutsche Forschungsgemeinschaft through the Sonderforschungsbereich 237 “Unordnung und große Fluktuationen”. One of us (A. Csordás) would like to acknowledge additional support by The Hungarian National Scientific Research Foundation under Grant number F4472.

## References

- [1] A. Macías, O. Obregón, and M. P. Ryan Jr., *Class. Quantum Grav.* **4**, 1477 (1987)
- [2] P. D. D'Eath and D. I. Hughes, *Phys. Lett.* **B214**, 498 (1988); *Nucl. Phys.* **B378**, 381 (1991)
- [3] J. Socorro, O. Obregón, and A. Macías, *Phys. Rev.* **D45**, 2026 (1992); O. Obregón, J. Socorro and J. Benitez, *ibid* **47**, 4471 (1993)
- [4] R. Graham *Phys. Rev. Lett* **67**, 1381 (1991); *Phys. Lett.* **B277**, 393 (1992); J. Bene and R. Graham, *Phys. Rev.* **D49**, 799 (1994)
- [5] P. D. D'Eath, S. W. Hawking and O. Obregón, *Phys. Lett.* **B300**, 44 (1993)
- [6] P. D. D'Eath, *Phys. Rev.* **D48**, 713 (1993)
- [7] R. Graham *Phys. Rev.* **D48**, 1602 (1993)
- [8] M. Asano, M. Tanimoto, and N. Yoshino, *Phys. Lett.* **B314**, 303 (1993)
- [9] R. Capovilla and J. Guven (preprint 1994, gr-qc/9402025)
- [10] P. D. D'Eath, *Phys. Lett.* **B320**, 12 (1994)
- [11] A. D. Y. Cheng, P. D. D'Eath and P. R. L. V. Moniz, *Phys. Rev.* **D49**, 5246 (1994); (preprint 1994, gr-qc/9406047)
- [12] R. Capovilla, and O. Obregón, *Phys. Rev.* **D49**, 6562 (1994)
- [13] R. Graham and H. Luckock, *Phys. Rev.* **D49**, R4981 (1994)
- [14] S. M. Carroll, D. Z. Freedman, M E. Ortiz, and D. N. Page, *Nucl. Phys.* **B423**, 661 (1994)
- [15] G. F. R. Ellis and M. A. H. McCallum, *Comm. Math. Phys.* **12**, 108 (1960)
- [16] J. Wess and J. Bagger, “Supersymmetry and Supergravity”, (Princeton University Press, Princeton 1972)
- [17] P. D. D'Eath, *Phys. Rev.* **D29**, 2199 (1984)
- [18] M. P. Ryan Jr. and L. C. Shepley, “Homogeneous Relativistic Cosmologies” (Princeton University Press, Princeton 1975)
- [19] A. Csordás and R. Graham, to appear in *Phys. Rev. Lett.*, April 1995

[20] We note that this choice of ordering is *opposite* to the one adopted in [19] but just as legitimate. Correspondingly the generator algebra below and the form of eqs. (12), (14) also slightly differ from [19] but both versions are fully consistent with each other. The choice of ordering adopted here is motivated by the simplification it brings to eq. (18) in the application given towards the end of this lecture.